10mm clear area for Schwabe pharmacoding

Each film-coated tablet contains 124mg of lactose and 36mg of liquid glucose. Each pack contains 30 film-coated tablets.

Registration holder for this product

Schwabe Pharma (UK) Ltd, Alexander House, Mere Park, Dedmere Road, Marlow, Bucks, SL7 1FX

Manufacturer of this product

Wiewelhove GmbH, Gildestrasse 39, 49477 Ibbenbüren, Germany Traditional herbal registration number: THR 23056/0002

If you would like further information about this product or if this leaflet is hard to see or read, please contact: Healthspan Ltd, PO Box 64, Guernsey, GY1 1BT.

THR 23056/0002

certification mark

THR

Telephone: 0800 73 123 77.

Email: herbalinfo@healthspan.co.uk

This leaflet was revised in March 2018.

Patient Information Leaflet

Healthspan Agnus Castus Premenstrual Relief

Agnus Castus Fruit Extract 4mg

Please read this leaflet carefully before you start taking these tablets.

It contains some important information about Healthspan Agnus Castus Premenstrual Relief.

Keep this leaflet with the tablets.

You may want to read it again or show it to your doctor, pharmacist or qualified healthcare practitioner.

What is in this leaflet

:	What this product is and what it is used for	page 1
:	Before you take this product	page 2
:	How to take this product	page 2-3
:	Possible side-effects	page 3-4
:	How to store this product	page 4
	Further information	nage 4-5

1: What this product is and what it is used for

This product is a traditional herbal medicinal product containing Agnus Castus fruit. Each film-coated tablet contains 4mg of extract (as dry extract) from Agnus Castus fruit (Vitex agnus castus L.) (7-13:1) (equivalent to 28-52mg of Agnus Castus). Extraction solvent: Ethanol 60% v/v.

Healthspan Agnus Castus Premenstrual Relief is a traditional herbal medicinal product used to help relieve symptoms of premenstrual syndrome such as irritability, mood swings, breast tenderness, bloating and menstrual cramps. This usage is based on traditional use only.

DO NOT TAKE this product if you are:
pregnant or trying to become pregnant

- breastfeeding
- allergic to any of the ingredients (see section 6)
- suffering from a pituitary disorder
- under 18 years of age

Tell your doctor before taking this product if you:

- have an intolerance to some sugars (see section 6)
- have had a pituitary disorder in the past
- suffer from, or have suffered from, an oestrogen-sensitive cancer
- are taking any of the following: dopamine receptor antagonist or agonist type medicine, oestrogens or antioestrogens or any other medicines including any bought without a prescription.

3: How to take this product

For adult women experiencing premenstrual symptoms, take 1 tablet daily

Try to take the tablets at the same time each day, either in the morning or evening. Swallow the tablets whole with some water or other liquid. Do not chew the tablets. You can take the tablets with or without food. The maximum beneficial effect of this product may take up to 3 months to develop in some women.

Do not exceed the stated dose.

Do not give to children or adolescents under 18 years old.

If symptoms worsen, or persist, or do not improve after 3 months, a doctor or qualified practitioner should be consulted.

If you take too much of this product (overdose)

If you take more than the recommended dose, speak to a doctor, pharmacist or qualified healthcare practitioner and take this leaflet with you.

If you forget to take this product

Do not take a double dose to make up for the missed dose. Continue to take your usual dose at the usual time, it does not matter if you have missed a dose.

If you have any questions, or are unsure about anything, please ask your doctor, pharmacist or qualified healthcare practitioner.

4: Possible side-effects

Like all medicines this product can have side-effects. Side-effects that have been reported are listed below.

Side-effects

 severe allergic reactions such as swelling of the face, breathlessness or difficulty swallowing. If you experience these effects, seek medical advice immediately.

Other side-effects reported

- nausea or feeling sick
- · abdominal pain
- menstrual disorders
- acne
- dizziness
- headache
- allergic skin reactions itching and/or rash of the skin

The frequency of these effects occurring is not known. If these persist for more than a few days, or become troublesome, stop taking this product. These side-effects are often only temporary. Stop taking this product immediately if you experience any allergic skin reaction.

Other side-effects

Tell your doctor or pharmacist if you notice any other side-effect not mentioned in this leaflet.

Reporting of side-effects

If you get any side-effects, talk to your doctor, pharmacist or qualified healthcare practitioner. This includes any possible side-effects not listed in this leaflet. You can also report side-effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side-effects you can help provide more information on the safety of this medicine.

5: How to store this product

Do not use your tablets after the expiry date. Return any out-of-date tablets to your pharmacist who will dispose of them for you. The expiry date is printed on the box and the blister pack.

This medicinal product does not require any special storage conditions.

Keep the tablets out of the reach and sight of children.

Keep your tablets in the blister pack until it is time to take them.

6: Further information

Each film-coated tablet contains 4mg of extract (as dry extract) from Agnus Castus fruit (*Vitex agnus castus* L.) (7-13:1) (equivalent to 28-52mg of Agnus Castus). Extraction solvent: Ethanol 60% v/v.

This product also contains the following ingredients:

Tablet core: Liquid glucose, silica colloidal anhydrous, lactose monohydrate, magnesium stearate, maize starch, cellulose microcrystalline, sodium starch glycolate (type A).

Film-coating: Lactose monohydrate, hypromellose, macrogol 4000, titanium dioxide (E171), iron (III)-oxide (E172).

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this product.